

## WEST Search History

DATE: Tuesday, June 24, 2003

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>			
L2	L1 and (transgen\$ or knockout)	5	L2
L1	mageko	5	L1

END OF SEARCH HISTORY

BEST AVAILABLE COPY

\$%^STN;HighlightOn= \*\*\*;HighlightOff=\*\*\* ;

Welcome to STN International! Enter x:x

LOGINID:ssspta1633cxq

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 Jun 03 New e-mail delivery for search results now available  
NEWS 4 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN  
NEWS 5 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE)  
now available on STN  
NEWS 6 Aug 28 Sequence searching in REGISTRY enhanced  
NEWS 7 Sep 03 JAPIO has been reloaded and enhanced  
NEWS 8 Sep 16 Experimental properties added to the REGISTRY file  
NEWS 9 Sep 16 CA Section Thesaurus available in CAPLUS and CA  
NEWS 10 Oct 01 CASREACT Enriched with Reactions from 1907 to 1985  
NEWS 11 Oct 24 BEILSTEIN adds new search fields  
NEWS 12 Oct 24 Nutraceuticals International (NUTRACEUT) now available on  
STN  
NEWS 13 Nov 18 DKILIT has been renamed APOLLIT  
NEWS 14 Nov 25 More calculated properties added to REGISTRY  
NEWS 15 Dec 04 CSA files on STN  
NEWS 16 Dec 17 PCTFULL now covers WP/PCT Applications from 1978 to date  
NEWS 17 Dec 17 TOXCENTER enhanced with additional content  
NEWS 18 Dec 17 Adis Clinical Trials Insight now available on STN  
NEWS 19 Jan 29 Simultaneous left and right truncation added to COMPENDEX,  
ENERGY, INSPEC  
NEWS 20 Feb 13 CANCERLIT is no longer being updated  
NEWS 21 Feb 24 METADEX enhancements  
NEWS 22 Feb 24 PCTGEN now available on STN  
NEWS 23 Feb 24 TEMA now available on STN  
NEWS 24 Feb 26 NTIS now allows simultaneous left and right truncation  
NEWS 25 Feb 26 PCTFULL now contains images  
NEWS 26 Mar 04 SDI PACKAGE for monthly delivery of multife SDI results  
NEWS 27 Mar 20 EVENTLINE will be removed from STN  
NEWS 28 Mar 24 PATDPFULL now available on STN  
NEWS 29 Mar 24 Additional information for trade-named substances without  
structures available in REGISTRY  
NEWS 30 Apr 11 Display formats in DGENE enhanced  
NEWS 31 Apr 14 MEDLINE Reload  
NEWS 32 Apr 17 Polymer searching in REGISTRY enhanced  
NEWS 33 Jun 13 Indexing from 1947 to 1956 added to records in CA/CAPLUS  
NEWS 34 Apr 21 New current-awareness alert (SDI) frequency in  
WPIDSWPINDEXPWPIX  
NEWS 35 Apr 28 RDISCLOSURE now available on STN  
NEWS 36 May 05 Pharmacokinetic information and systematic chemical names  
added to PHAR  
NEWS 37 May 15 MEDLINE file segment of TOXCENTER reloaded  
NEWS 38 May 15 Supporter information for ENCOMPPAT and ENCOMPLIT  
updated  
NEWS 39 May 16 CHEMREACT will be removed from STN  
NEWS 40 May 19 Simultaneous left and right truncation added to WSCA  
NEWS 41 May 19 RAPRA enhanced with new search field, simultaneous left and  
right truncation  
NEWS 42 Jun 06 Simultaneous left and right truncation added to CBNB  
NEWS 43 Jun 06 PASCAL enhanced with additional data  
NEWS 44 Jun 20 2003 edition of the FSTA Thesaurus is now available  
  
NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a,  
CURRENT  
MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),  
AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 15:20:03 ON 24 JUN 2003

=> FIL BIOSIS EMBASE CAPLUS

COST IN U.S. DOLLARS SINCE FILE TOTAL  
ENTRY SESSION

FULL ESTIMATED COST 0.21 0.21

FILE 'BIOSIS' ENTERED AT 15:20:10 ON 24 JUN 2003  
COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 15:20:10 ON 24 JUN 2003  
COPYRIGHT (C) 2003 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 15:20:10 ON 24 JUN 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

=> s mageko  
L1 0 MAGEKO

=> s transgen? or knockout  
L2 187730 TRANSGEN? OR KNOCKOUT

=> s l2 (3a)(mouse or mice)  
L3 99284 L2 (3A)(MOUSE OR MICE)

=> s l1 or knock-in or knockin or knock in  
L4 12923 L1 OR KNOCK-IN OR KNOCKIN OR KNOCK IN

=> s l2 or l4  
L5 197392 L2 OR L4

=> s l5 (3a) (mouse or mice)  
L6 102656 L5 (3A) (MOUSE OR MICE)

=> s l6 and VP16 and tetracycline repressor and GFP and VDE and neo  
L7 0 L6 AND VP16 AND TETRACYCLINE REPRESSOR AND GFP AND  
VDE AND NEO

=> s l6 and VP16 and tetracycline repressor and GFP and VDE  
L8 0 L6 AND VP16 AND TETRACYCLINE REPRESSOR AND GFP AND  
VDE

=> s l6 and VP16 and tetracycline repressor and GFP  
L9 0 L6 AND VP16 AND TETRACYCLINE REPRESSOR AND GFP

=> s l6 and VP16 and tetracycline repressor  
L10 9 L6 AND VP16 AND TETRACYCLINE REPRESSOR

=> dup rem l10  
PROCESSING COMPLETED FOR L10  
L11 5 DUP REM L10 (4 DUPLICATES REMOVED)

=> d bib abs 1-  
YOU HAVE REQUESTED DATA FROM 5 ANSWERS - CONTINUE? Y(N):y

L11 ANSWER 1 OF 5 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS  
INC.

AN 2001:311638 BIOSIS

DN PREV200100311638

TI C/EBPepsilon mediates myeloid differentiation and is regulated by the  
CCAAT displacement protein (CDP/cut) in 32Dc13 cells.

AU Khanna-Gupta, Arati (1); Zibello, Theresa (1); Lekstrom-Himes, Julie;  
Berliner, Nancy (1)

CS (1) Internal Medicine/Hematology, Yale University School of Medicine, New  
Haven, CT USA

SO Blood, (November 16, 2000) Vol: 96, No. 11 Part 1, pp. 282a. print.

Meeting Info.: 42nd Annual Meeting of the American Society of Hematology  
San Francisco, California, USA December 01-05, 2000 American Society of  
Hematology

ISSN: 0006-4971.

DT Conference

LA English

SL English

AB The role of C/EBPepsilon has been confirmed in \*\*\*knockout\*\*\*  
\*\*\*mice\*\*\*, which develop normally but fail to produce functional  
neutrophils. These mice die of opportunistic infections at 3-5 months of  
age. Neutrophils from C/EBPepsilon -/- mice have morphological and  
biochemical features similar to patients with neutrophil-specific  
secondary granule deficiency (SGD), a congenital disorder characterized by  
frequent bacterial infections and neutrophils that lack secondary granule  
proteins (SGP). Analysis of one SGD patient revealed a 5-basepair deletion  
in the C/EBPepsilon gene, resulting in an inactive truncated protein. Lack  
of functional C/EBPepsilon is postulated to cause this patient's disease.  
We previously demonstrated that overexpression of the transcriptional  
repressor CCAAT displacement protein (CDP/cut) in 32Dc13 cells suppresses  
SGP gene expression. This phenotype resembles that of C/EBPepsilon -/- mice and  
SGD patients. We therefore hypothesized that C/EBPepsilon was regulated by  
CDP/cut. We expressed wild type and mutant SGD C/EBPepsilon genes in  
32Dc13/Tet cells, which constitutively express a \*\*\*Tetracycline\*\*\*  
\*\*\*repressor\*\*\* (TetR)- \*\*\*VP16\*\*\* fusion protein. Stable  
transfection of 32D/Tet cells with a C/EBPepsilon cDNA cloned downstream  
of three TetR binding elements results in inducible C/EBPepsilon  
expression upon removal of Tet from the medium. Wild type but not mutant  
C/EBPepsilon expression in the 32D/Tet cells resulted in granulocytic  
morphology and lactoferrin (LF) gene expression. These data indicate that  
inducible C/EBPepsilon in 32D/Tet cells can drive granulocytic maturation.

BEST AVAILABLE COPY

Since SGP gene expression is dependent on C/EBPepsilon, and that expression is abrogated in 32Dcl3 CDP/cut overexpressing cells, we hypothesized that lack of SGP gene expression may reflect altered levels of C/EBPepsilon. Northern blot analysis of G-CSF-induced CDP/cut overexpressing 32Dcl3 cells revealed absence of C/EBPepsilon mRNA. Levels of C/EBPalpha transcript were unaffected. We conclude that repression of SGP gene expression in 32Dcl3/CDP/cut cells is mediated in two ways. The first is a direct interaction of CDP/cut with the promoters of the SGP genes, an observation we previously validated with the LF promoter. The second is an interaction between CDP/cut and C/EBPepsilon. We have isolated the C/EBPepsilon promoter and have identified a putative CDP/cut binding site. We have evidence for a role of this site in C/EBPepsilon expression and hence in SGP expression during neutrophil maturation.

L11 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2003 ACS

AN 1996:746323 CAPLUS

DN 126:15526

TI Glucose-responsive, insulin-producing transgenic pancreatic .beta.-cells with proliferation regulated by tetracycline

IN Efrat, Shimon

PA Albert Einstein College of Medicine of Yeshiva University, USA

SO PCT Int. Appl., 33 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 0631242	A1	19961010	WO 1996-US4792	19960403
W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2217652	AA	19961010	CA 1996-2217652	19960403
AU 9655375	A1	19961023	AU 1996-55375	19960403
AU 720662	B2	20000608		
EP 822834	A1	19980211	EP 1996-912616	19960403
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 11505411	T2	19990521	JP 1996-530530	19960403
US 6114599	A	20000905	US 1998-44297	19980319
US 6242254	B1	20010605	US 2000-492905	20000127
PRAI US 1995-418416	A	19950407		
WO 1996-US4792	W	19960403		
US 1998-44297	A1	19980319		
AB Glucose-regulated insulin producing pancreatic .beta.-cells whose proliferation is controlled by tetracyclines are described for use in the treatment of diabetes. Proliferation is controlled by a fusion protein of the ***tetracycline*** ***repressor*** tetR and ***VP16*** to regulate expression of an SV40 T antigen gene under control of a tet operator. The gene for the fusion protein is under control an insulin-responsive promoter. An animal carrying both constructs is prep. by crossing animals transformed with one of the constructs and .beta.-cells carrying the both constructs are selected in vitro. The construction of these cells in mice is demonstrated.				

L11 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2003 ACS

AN 1997:51843 CAPLUS

DN 126:100260

TI Fusion proteins of the \*\*\*tetracycline\*\*\* \*\*\*repressor\*\*\* for use in tetracycline regulation of gene expression in eukaryotes

IN Bujard, Hermann; Gossen, Manfred; Hillen, Wolfgang; Helbl, Vera; Schnappinger, Dirk

PA BASF A.-G., Germany; Knoll Aktiengesellschaft

SO U.S., 62 pp., Cont.-in-part of U.S. Ser. No. 383,754.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 12

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI US 5589362	A	19961231	US 1995-485971	19950607
US 5464758	A	19951107	US 1993-76726	19930614
US 5650298	A	19970722	US 1994-260452	19940614
US 5654168	A	19970805	US 1994-275876	19940715
US 5789156	A	19980804	US 1995-383754	19950203
WO 9640892	A1	19961219	WO 1996-US9049	19960606
W: CA, JP				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
PRAI US 1993-76327	B2	19930614		
US 1993-76726	A2	19930614		
US 1994-260452	A2	19940614		
US 1994-270637	B2	19940701		
US 1994-275876	A2	19940715		
US 1995-383754	A2	19950203		
US 1995-485971	A	19950607		
AB Fusion proteins of amino acid-substituted tet repressors and transcription factors that bind class B tet operators that can be used in tetracycline regulation of expression of foreign genes in eukaryotes. Genes encoding these proteins are also described. The tet operators also have nucleotide				

substitutions in one or two of the 3'-bases (+4 or +6). A pool of multiply mutant tet repressor genes was generated by bisulfite mutagenesis of the tetR gene and mutants with a reverse regulation phenotype (induction of gene expression by tetracyclines rather than repression) were identified using a galK/lacZ/tet operator reporter system. Fusion proteins of the N-terminal regions of these proteins and herpes simplex \*\*\*VP16\*\*\* were prep. by std. methods. Their efficacy was tested in a reporter gene system using the CMV promoter and a heptameric tet operator to regulate expression of a luciferase reporter in HR-5 cells. Doxycycline induced gene expression by 237-1660-fold and two genes under the control of tet operators could be induced coordinately. Fusion proteins of silencer domains, e.g. Krueppel or v-erbA proteins, are described for use as repressors. A combinatorial anal. of amino acid-substituted analogs of the repressor and base-substituted analogs of the operator was undertaken to find combinations showing the most effective induction or repression.

L11 ANSWER 4 OF 5 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE 1

AN 1996:537416 BIOSIS

DN PREV199699259772

TI Temporal control of the Cre recombinase in \*\*\*transgenic\*\*\*

\*\*\*mice\*\*\* by a tetracycline responsive promoter.

AU St-Onge, Luc; Furth, Priscilla A.; Gruss, Peter (1)

CS (1) Dep. Mol. Cell. Biol., Max-Planck-Inst. Biophys. Chem., Am Fassberg, 37018 Goettingen Germany

SO Nucleic Acids Research, (1996) Vol. 24, No. 19, pp. 3875-3877.

ISSN: 0305-1048.

DT Article

LA English

AB Gene-targeted mice derived from embryonic stem cells are a useful tool to study gene function during development. However, if the mutation is embryonic lethal and the gene is deleted from the onset of development, later functions in adult animals cannot be studied. Recently, the bacterial Cre-loxP site-specific recombination system has successfully been used in transgenic animals to produce tissue-specific and temporal deletions (Gu et al. (1993) Cell, 73, 1155-1164; Gu et al. (1994) Science, 265, 103-106; Kuhn et al. (1995) Science, 269, 1427-1429). We have evaluated the tetracycline responsive binary system (Gossen and Bujard (1992) Proc. Natl. Acad. Sci. USA, 89, 5547-5551) for its ability to transiently express the Cre recombinase in \*\*\*transgenic\*\*\* \*\*\*mice\*\*\*. In this system, a transactivator fusion protein composed of the \*\*\*tetracycline\*\*\* \*\*\*repressor\*\*\* (tetR) and the acidic domain of the herpes simplex viral protein 16 ( \*\*\*VP16\*\*\* ) can regulate the expression of the Cre gene from a promoter containing tet-operator (tetO) sequences. In the absence of tetracycline, the Cre gene is expressed and will induce site-specific recombination between two loxP sites. In the presence of tetracycline, the Cre gene will not be expressed and recombination will not occur.

L11 ANSWER 5 OF 5 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE 2

AN 1995:361280 BIOSIS

DN PREV199598375580

TI A modified tetracycline-regulated system provides autoregulatory,

inducible gene expression in cultured cells and \*\*\*transgenic\*\*\*

\*\*\*mice\*\*\*

AU Shockett, Penny; Difilippantonio, Michael; Hellman, Nathan; Schatz, David G. (1)

CS (1) Howard Hughes Med. Inst., Yale Univ. Sch. Med., New Haven, CT 06510 USA

SO Proceedings of the National Academy of Sciences of the United States of America, (1995) Vol. 92, No. 14, pp. 6522-6526.

ISSN: 0027-8424.

DT Article

LA English

AB A system for tetracycline-regulated inducible gene expression was described recently which relies on constitutive expression of a tetracycline-controlled transactivator (tTA) fusion protein combining the \*\*\*tetracycline\*\*\* \*\*\*repressor\*\*\* and the transcriptional activation domain of \*\*\*VP16\*\*\* (Gossen, M. & Bujard, H. (1992) Proc. Natl. Acad. Sci. US. A 89, 5547-5551). This system yielded only low levels of transactivator protein, probably because tTA is toxic. To avoid this difficulty, we placed the tTA gene under the control of the inducible promoter to which tTA binds, making expression of tTA itself inducible and autoregulatory. When used to drive expression of the recombination activating genes 1 and 2 (RAG-1 and RAG-2), the autoregulatory system yielded both substantially higher levels of variable (diversity) joining (V(D)J) recombination activity (70-fold on average) and inducible expression in a much larger fraction of transfected cells (autoregulatory, 90%, vs. constitutive, 18%). In addition, this system allowed the creation of \*\*\*transgenic\*\*\* \*\*\*mice\*\*\* in which expression of a luciferase transgene was inducible tens to hundreds of times the basal levels in most tissues examined. Induced levels of expression were highest in thymus and lung and appear to be substantially higher than in previously reported inducible luciferase \*\*\*transgenic\*\*\* \*\*\*mice\*\*\* created with the constitutive system. With the modified system, inducible transactivator mRNA and protein were easily detected in cell lines by RNA and Western blotting, and transactivator mRNA was detected by RNA blotting in some tissues of \*\*\*transgenic\*\*\* \*\*\*mice\*\*\*. This autoregulatory system represents an improved strategy for tetracycline-regulated gene expression both in cultured cells and in transgenic animals.

=> FIL STNGUIDE  
COST IN U.S. DOLLARS      SINCE FILE    TOTAL  
ENTRY    SESSION  
FULL ESTIMATED COST      78.81    79.02

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)    SINCE FILE  
TOTAL

ENTRY    SESSION  
CA SUBSCRIBER PRICE      -1.30    -1.30

FILE 'STNGUIDE' ENTERED AT 15:34:25 ON 24 JUN 2003  
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM  
KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.  
LAST RELOADED: Jun 20, 2003 (20030620/UP).

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS      SINCE FILE    TOTAL  
ENTRY    SESSION  
FULL ESTIMATED COST      0.24    79.26

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)    SINCE FILE  
TOTAL

ENTRY    SESSION  
CA SUBSCRIBER PRICE      0.00    -1.30

STN INTERNATIONAL LOGOFF AT 15:36:35 ON 24 JUN 2003